



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

CALCULUS.

220. Proposed by C. N. SCHMALL, College of the City of New York, New York City.

To determine the least polygon of n sides that can be described about a given circle.

Solution by the PROPOSER.

Let $\phi_1, \phi_2, \dots, \phi_n$ be the successive angles contained between the lines l_1, l_2, \dots , drawn from the center to the vertices of the polygon, and the radii (r) drawn to the points of contact of the sides. The area of the right triangle whose angle at the center is ϕ_1 , will be

$$\Delta = \frac{1}{2} r l_1 \sin \phi_1 = \frac{1}{2} r \cdot r \sec \phi_1 \cdot \sin \phi_1 = \frac{r^2}{2} \tan \phi_1.$$

Hence the entire area of the polygon is

$$\Sigma \Delta = \frac{r^2}{2} [\tan \phi_1 + \tan \phi_2 + \dots + \tan \phi_n].$$

But $\tan \phi_n = \tan[2\pi - (\phi_1 + \phi_2 + \dots + \phi_{n-1})] = \tan[2\pi - \theta]$, where $\theta = \phi_1 + \phi_2 + \dots + \phi_{n-1}$. Thus $u = \tan \phi_1 + \tan \phi_2 + \dots + \tan(2\pi - \theta)$ is to be rendered a minimum. Differentiating partially with respect to ϕ_1 , we obtain

$$\frac{\partial u}{\partial \phi_1} = \sec^2 \phi_1 - \sec^2(2\pi - \theta) = 0.$$

Hence $\phi_1 = 2\pi - \theta = \phi_n$.

In like manner we may show that any angle equals the one preceding it. Hence the minimum polygon is regular.

Also solved by G. W. Greenwood, and J. Scheffer.

DIOPHANTINE ANALYSIS.

136. Proposed by A. H. HOLMES, Brunswick, Maine.

Given $7x^2 - 111 = y^2$. Required a value for y greater than unity which shall be a prime integer.

Solution by J. SCHEFFER, A. M., Hagerstown, Md.

Let $y = q$, $x = p$ be two values that satisfy the equation $y^2 - Nx^2 = -a$, and $y = n$, $x = m$ two values that satisfy the equation $y^2 - Nx^2 = 1$. Then we evidently have $y^2 - Nx^2 = (q^2 - Np^2)(n^2 - Nm^2) = n^2q^2 + N^2m^2p^2 - N(m^2q^2 + n^2p^2) = (nq \pm Npm)^2 - N(mq \pm np)^2$. Therefore, we can put $y = nq \pm Npm$, $x = mq \pm np$. Substituting numerical values we have, since $y = 8$, $x = 3$ satisfy the equation $y^2 - 7x^2 = 1$,